

Holomorphic Morse Inequalities on Covering Manifolds

Radu Todor, Ionuț Chiose

Abstract

The goal of this paper is to generalize Demailly's asymptotic holomorphic Morse inequalities to the case of a covering manifold of a compact manifold. We shall obtain estimates which involve Atiyah's "normalized dimension" of the square integrable harmonic spaces. The techniques used are those of Shubin who gave a proof for the usual Morse inequalities in the presence of a group action relying on Witten ideas. As a consequence we obtain estimates for the dimension of the square integrable holomorphic sections of the pull-back of a line bundle on the base manifold under some mild hypothesis for the curvature.

1 Introduction

To emphasize the meaning of our result let us consider a compact projective manifold X , F an ample line bundle on X , $\tilde{X}$ the universal covering of X and $\tilde{F}$ the pull-back of F on $\tilde{X}$. Then Kollár's Theorem 6.4 from [5] shows that

$$\dim_{\pi_1(X)} H_{(2)}^0(\tilde{X}, K_{\tilde{X}} \otimes \tilde{F}) = \dim H^0(X, K_X \otimes F)$$

where we denote as usual K_N the canonical bundle of a manifold N . It follows from Theorem 0.1 in Demailly [4] that $\dim_{\pi_1(X)} H_{(2)}^0(\tilde{X}, K_{\tilde{X}} \otimes \tilde{F}^k)$ has polynomial growth of order the dimension of X as $k \rightarrow \infty$. We shall generalize this result to the case of a complex analytic manifold M on which a discrete group Γ acts freely and properly discontinuous such that $X = M/\Gamma$ is compact and carries a line bundle F satisfying Demailly's condition:

$$\int_{X(\leq 1)} (ic(F))^n > 0 \quad (1)$$

Let us mention that holomorphic Morse inequalities on non-compact manifolds have been obtained before on q -concave and q -convex manifolds (see Marinescu [6] and Bouche [3]).

The cohomology groups of M are usually infinite-dimensional and we cannot use the usual dimension. The dimension we shall use is the Γ -dimension introduced by Atiyah in [2]. Let us define the Γ -dimension of certain subspaces of $L^2(M, \tilde{E})$, where $E \rightarrow X$ is a hermitian vector bundle, $\tilde{E} \rightarrow M$ is the pull-back of E , X (and hence M) is endowed with a Riemannian metric and

$$L^2(M, \tilde{E}) = \{s : M \rightarrow \tilde{E} \mid s \text{ is a measurable section, } \int_M |s|^2 dV < \infty\}$$

Let G be a closed subspace in $L^2(M, \tilde{E})$ such that $L_\gamma G \subset G$ where L_γ is the action of Γ on $L^2(M, \tilde{E})$. Let U be a fundamental domain for the action of Γ and $(\varphi_m)_m$

an orthonormal base for the Hilbert space G . Then one defines

$$\dim_{\Gamma} G := \sum_m \int_U |\varphi_m(x)|^2 dV(x) \quad (2)$$

It can be shown that the definition of $\dim_{\Gamma} G$ does not depend on the orthonormal base $(\varphi_m)_m$ or on the fundamental domain U (see Atiyah [2]).

Let M be a complex analytic manifold of complex dimension n on which a discrete group Γ acts freely and properly discontinuous such that $X = M/\Gamma$ is compact. Since M/Γ is compact one can easily see that the pull-back of a hermitian metric on X is a complete metric on M which we consider fixed from now on. Let E be a hermitian holomorphic vector bundle on X and $\tilde{E} = \pi^* E$ its pull-back, where $\pi : M \rightarrow X$ is the projection.

Let

$$\bar{\partial}_q : C_{0,q}^{\infty}(M, \tilde{E}) \rightarrow C_{0,q+1}^{\infty}(M, \tilde{E})$$

be the well-known Cauchy–Riemann operator and

$$\delta_q : C_{0,q+1}^{\infty}(M, \tilde{E}) \rightarrow C_{0,q}^{\infty}(M, \tilde{E})$$

the formal adjoint of $\bar{\partial}_q$. Then $\Delta_q'' = \bar{\partial}_{q-1} \delta_{q-1} + \delta_q \bar{\partial}_q$ is an elliptic differential operator.

Let $\bar{\partial}_q : L_{0,q}^2(M, \tilde{E}) \rightarrow L_{0,q+1}^2(M, \tilde{E})$ be the weak maximal extension of $\bar{\partial}_q$ and likewise we denote by the same letter the weak maximal extensions of δ_q and Δ_q'' . Let us denote by $N^q(\bar{\partial})$ the kernel of $\bar{\partial}_q$, by $R^{q-1}(\bar{\partial})$ the range of $\bar{\partial}_{q-1}$, $N^q(\delta)$ the kernel of $\delta_{q-1} : L_{0,q}^2(M, \tilde{E}) \rightarrow L_{0,q-1}^2(M, \tilde{E})$ and by $N^q(\Delta'')$ the kernel of $\Delta_q'' : L_{0,q}^2(M, \tilde{E}) \rightarrow L_{0,q}^2(M, \tilde{E})$.

By basic results of Andreotti and Vesentini [1] the hilbertian adjoint of $\bar{\partial}$ coincides with δ , Δ_q'' is self-adjoint and

$$\mathcal{H}_{(2)}^q(M, \tilde{E}) := N^q(\Delta'') = N^q(\delta) \cap N^q(\bar{\partial})$$

where the first equality is the definition of the space of L^2 harmonic forms.

Let E and F be hermitian holomorphic fibre bundles on X of rank 1 and r respectively, $\tilde{F} = \pi^* F$, $\tilde{E} = \pi^* E$. Let us denote $D = D' + \bar{\partial}$ the canonical connection of E and $c(E) = D^2 = D' \bar{\partial} + \bar{\partial} D'$ its curvature form. Also, let

$$X(q) = \{x \in X \mid ic(E) \text{ has } q \text{ negative eigenvalues and } n - q \text{ positive eigenvalues}\}$$

and

$$X(\leq q) = X(0) \cup X(1) \cup \dots \cup X(q).$$

The main theorem of this paper is:

Theorem 1.1. *As $k \rightarrow \infty$, the following inequalities hold for every $q = 0, 1, \dots, n$:*

i) *The weak Morse inequalities:*

$$\dim_{\Gamma} \mathcal{H}_{(2)}^q(M, \tilde{E}^k \otimes \tilde{F}) \leq r \frac{k^n}{n!} \int_{X(q)} (-1)^q \left(\frac{i}{2\pi} c(E) \right)^n + o(k^n).$$

ii) *The strong Morse inequalities:*

$$\sum_{j=0}^q (-1)^{q-j} \dim_{\Gamma} \mathcal{H}_{(2)}^j(M, \tilde{E}^k \otimes \tilde{F}) \leq r \frac{k^n}{n!} \int_{X(\leq q)} (-1)^q \left(\frac{i}{2\pi} c(E) \right)^n + o(k^n).$$

iii) *The asymptotic Riemann-Roch formula:*

$$\sum_{j=0}^n (-1)^j \dim_{\Gamma} \mathcal{H}_{(2)}^j(M, \tilde{E}^k \otimes \tilde{F}) = r \frac{k^n}{n!} \int_X \left(\frac{i}{2\pi} c(E) \right)^n + o(k^n).$$

It follows easily

Corollary 1.2. *Let E and F be as above and suppose that E satisfies (1). Then the space of L^2 holomorphic sections satisfies*

$$\dim_{\Gamma} H_{(2)}^0(M, \tilde{E}^k \otimes \tilde{F}) \approx k^n$$

as $k \rightarrow \infty$. In particular the usual dimension of the space of L^2 holomorphic sections of E^k has the same cardinal as $|\Gamma|$ for large k .

This generalizes the result for the covering of a projective manifold by T. Napier [7].

We wish to express our gratitude to Professor V. Iftimie and Dr. G. Marinescu for their pertinent suggestions and for the support we have received.

2 Γ -dimension and Estimates

Let M be a real Riemann manifold of dimension n , Γ a discrete group acting freely and properly discontinuous on M such that $X = M/\Gamma$ is compact, $F \rightarrow X$ a hermitian vector bundle of rank r and $\tilde{F} \rightarrow M$ is the pull-back of F . Let U be a fundamental domain for the action of Γ . We identify $L^2(M, \tilde{F}) \cong L^2\Gamma \otimes L^2(U, \tilde{F}) \cong L^2\Gamma \otimes L^2(X, F)$. Let us consider $\mathcal{A}_{\Gamma}$ the von Neumann algebra of bounded operators on $L^2(M, \tilde{F})$ which commute with Γ .

If $A \in \mathcal{A}_{\Gamma}$, then let $K_A \in \mathcal{D}'(M \times M, \tilde{F} \otimes_{M \times M} \tilde{F})$ its kernel. As A is Γ -invariant, it follows that $K_A \in \mathcal{D}'(M \times M/\Gamma, \tilde{F} \otimes_{M \times M} \tilde{F}/\Gamma)$ where the action of Γ on $M \times M$ is $(x, y) \rightarrow (\gamma x, \gamma y)$.

$A \in \mathcal{A}_{\Gamma}$ is said to be Γ -Hilbert-Schmidt if $K_A \in L^2(M \times M/\Gamma, \tilde{F} \otimes_{M \times M} \tilde{F}/\Gamma)$ and of Γ -trace class if $A = A_1 A_2$ with A_1, A_2 being Γ -Hilbert-Schmidt. If $A \in \mathcal{A}_{\Gamma}$ is of Γ -trace-class, one can define

$$\mathrm{Tr}_{\Gamma} A := \mathrm{Tr}(\varphi A \psi) \quad (3)$$

where $\varphi, \psi \in L_{comp}^{\infty}(M)$ such that $\sum_{\gamma \in \Gamma} (\varphi \psi) \circ \gamma = 1$. If $L \subset L^2(M, \tilde{F})$ is a closed, Γ -invariant subspace, that is L is a Γ -module, and P_L is the ortogonal projection onto L , then

$$\dim_{\Gamma} L := \mathrm{Tr}_{\Gamma} P_L \in [0, \infty] \quad (4)$$

This is in short the theory of Γ -traces. For more results see Atiyah [2] and Shubin [8]. We shall use the following three results; for the proofs see Shubin [8].

Proposition 2.1. *Let*

$$0 \rightarrow L_0 \rightarrow L_1 \rightarrow \dots \rightarrow L_q \rightarrow L_{q+1} \rightarrow \dots \rightarrow L_n \rightarrow 0$$

be a complex of Γ -modules (d_q commutes with the action of Γ and $d_{q+1}d_q = 0$). If $l_q = \dim_{\Gamma} L_q < \infty$ and $\bar{h}_q = \dim_{\Gamma} \bar{H}_q(L)$ where

$$\bar{H}_q(L) = N(d_q) / \overline{R(d_{q-1})}$$

then

$$\sum_{j=1}^q (-1)^{q-j} \bar{h}_j \leq \sum_{j=1}^q (-1)^{q-j} l_j \quad (5)$$

for every $q = 0, 1, \dots, n$ and for $q = n$ the inequality becomes equality.

Let $H = H^*$ be a linear operator in $L^2(M, \tilde{F})$ which commutes with the action of Γ , that is $E_\lambda \in \mathcal{A}_\Gamma$, where $(E_\lambda)_\lambda$ is the spectral family of H . Let us denote $N_\Gamma(\lambda, H) = \dim_\Gamma R(E_\lambda)$ and h the quadratic form of H .

Proposition 2.2. *If $H \geq 0$, then*

$$N_\Gamma(\lambda, H) = \sup\{\dim_\Gamma L \mid L \text{ is a } \Gamma\text{-module} \subset \text{Dom}(h), \\ h(f, f) \leq \lambda((f, f)), \forall f \in L\}. \quad (6)$$

Proposition 2.3. *If there is $T : L^2(M, \tilde{F}) \rightarrow L^2(M, \tilde{F})$ a Γ -endomorphism (i.e. T commutes with the action of Γ) such that $((H + T)f, f) \geq \mu((f, f))$, $f \in \text{Dom}(H)$ and $\text{rank}_\Gamma T = \dim_\Gamma \bar{R}(T) \leq p$, then*

$$N_\Gamma(\mu - \varepsilon, H) \leq p, \quad \forall \varepsilon > 0. \quad (7)$$

Let H be an elliptic differential operator, formally self-adjoint of order $2m$ on $\tilde{F}$, which commutes with the action of Γ . We shall denote by the same letter H the weak maximal extension of H . If H is strongly elliptic, then H is bounded from below (see Shubin[8]).

Theorem 2.4. *Let H_0 be the self-adjoint operator in $L^2(U, \tilde{F}|_U)$ defined by the restriction of H to U with Dirichlet boundary conditions. (H_0 is bounded from below and has compact resolvent). Then*

$$N_\Gamma(\lambda, H) \geq N(\lambda, H_0), \quad \forall \lambda \in \mathbb{R} \quad (8)$$

where $N(\lambda, H) = \dim R(F_\lambda)$ if (F_λ) is the spectral family of H_0 .

Proof. Let $(e_i)_i$ be an orthonormal basis of $L^2(U, \tilde{F})$ which consists of eigenfunctions of H_0 corresponding to the eigenvalues $(\lambda_i)_i$; if we let $\tilde{e}_i = 0$ on $M \setminus U$ and $\tilde{e}_i = e_i$ on U , then $\tilde{e}_i \in \text{Dom}(h)$ and $(L_\gamma \tilde{e}_i)_{i, \gamma}$ is an orthonormal basis of $L^2(M, \tilde{F})$ and $\tilde{e}_{i, \gamma} = L_\gamma \tilde{e}_i \in \text{Dom}(h)$. We have $h(\tilde{e}_{i, \gamma}, \tilde{e}_{i', \gamma'}) = \delta_{i, i'} \delta_{\gamma, \gamma'} \lambda_i$. Let Φ_λ^0 be the subspace spanned by $(\tilde{e}_i)_{\lambda_i \leq \lambda}$ and Φ_λ the closed subspace spanned by $(\tilde{e}_{i, \gamma})_{\lambda_i \leq \lambda}$. Then

$$\dim_\Gamma \Phi_\lambda = \sum ((P_{\Phi_\lambda} \tilde{e}_i, \tilde{e}_i)) = \sum_{\lambda_i \leq \lambda} ((\tilde{e}_i, \tilde{e}_i)) = \dim \Phi_\lambda^0 = N(\lambda, H_0).$$

If f is a linear combination of $\tilde{e}_{i, \gamma}$, $\lambda_i \leq \lambda$, then $h(f, f) \leq \lambda \|f\|^2$ and, as $\text{Dom}(h)$ is complete, we obtain that $\Phi_\lambda \subset \text{Dom}(h)$ and $h(f, f) \leq \lambda \|f\|^2$, $f \in \Phi_\lambda$. From Proposition 2.2 it follows that $N_\Gamma(\lambda, H) \geq N(\lambda, H_0)$. $\square$

Let $s > 0$, $U_s = \{x \in M \mid d(x, U) < s\}$ where d is the distance on M associated to the Riemann metric on M and $U_{s, \gamma} := \gamma U_s$. Let $\varphi^{(s)} \in C_0^\infty(M)$, $\varphi^{(s)} \geq 0$, $\varphi^{(s)} = 1$ on $\bar{U}$ and $\text{supp } \varphi^{(s)} \subset U_s$, $\varphi_\gamma^{(s)} = \varphi^{(s)} \circ \gamma^{-1}$. Put

$$C_\gamma^{(s)} = \frac{\varphi_\gamma^{(s)}}{\left(\sum_\gamma (\varphi_\gamma^{(s)})^2\right)^{\frac{1}{2}}} \in C_0^\infty(M)$$

so that $\sum_{\gamma \in \Gamma} (C_\gamma^{(s)})^2 = 1$. If $m = 1$ (that is H is of order 2) then

$$H = \sum_{\gamma \in \Gamma} C_\gamma^{(s)} H C_\gamma^{(s)} - \sum_{\gamma \in \Gamma} \sigma_0(H)(dC_\gamma^{(s)}) \quad (9)$$

where σ_0 is the principal symbol of H (see Shubin[8]).

Let us assume that X is a complex analytic manifold, $\pi : M \rightarrow X$, E and F hermitian holomorphic vector bundles on X , $\tilde{E} = \pi^* E$, $\tilde{F} = \pi^* F$. Let $\Delta_{k,q}''$ be the Laplace–Beltrami operator on $\Lambda^{0,q} T^* M \otimes \tilde{E}^k \otimes \tilde{F}$. If $s = k^{-\frac{1}{4}}$, $H = \frac{1}{k} \Delta_{k,q}''$ in (9) then it follows that there is a constant C such that

$$\frac{1}{k} \Delta_{k,q}'' \geq \sum_{\gamma \in \Gamma} \frac{1}{k} J_\gamma^{(k)} \Delta_{k,q}'' J_\gamma^{(k)} - \frac{C}{\sqrt{k}} \text{Id} \quad (10)$$

where $J_\gamma^{(k)} = C_\gamma^{(k^{-\frac{1}{4}})}$, $k \in \mathbb{N}^*$. We have used that $\sigma_0(\Delta'')(dJ) = |\bar{\partial} J|^2 \text{Id}$ if $J \in C^\infty(M, \mathbb{R})$.

Let us denote by $V = \Lambda^{0,q} T^* M \otimes \tilde{E}^k \otimes \tilde{F}$, $H = \frac{1}{k} \Delta_{k,q}''$ and $H_0^{(k)} = \frac{1}{k} \Delta_{k,q}''|_{U_{k^{-\frac{1}{4}}}}$ the operator defined in $L^2(U_{k^{-\frac{1}{4}}}, V)$ by the restriction of $\frac{1}{k} \Delta_{k,q}''$ to $U_{k^{-\frac{1}{4}}}$ with Dirichlet boundary conditions. Let $(E_\lambda^{(k)})_\lambda$ be the spectral family of $H_0^{(k)}$. In the sequel we fix λ and consider $M^{(k)}$ a real number such that $M^{(k)} \geq \lambda - \inf \text{spec}(H_0^{(k)})$, where $\text{spec}(H_0^{(k)})$ is the spectrum of $H_0^{(k)}$ to the effect that

$$H_0^{(k)} + M^{(k)} E_\lambda^{(k)} \geq \lambda \text{Id}.$$

Define

$$\begin{aligned} G_\gamma^{(k)} &: L^2(M, V) \rightarrow L^2(M, V) \\ G_\gamma^{(k)} &= J_\gamma^{(k)} L_\gamma M^{(k)} E_\lambda^{(k)} L_\gamma^{-1} J_\gamma^{(k)}, \end{aligned}$$

(i.e. we trunk the section over $U_{s,\gamma}$, transport it on U_s , apply the spectral projection and then send it back to $U_{s,\gamma}$) and

$$G^{(k)} = \sum_{\gamma \in \Gamma} G_\gamma^{(k)}.$$

We have

$$\begin{aligned} H + G^{(k)} &\geq \sum_{\gamma \in \Gamma} \left(J_\gamma^{(k)} H J_\gamma^{(k)} + J_\gamma^{(k)} L_\gamma M^{(k)} E_\lambda^{(k)} L_\gamma^{-1} J_\gamma^{(k)} \right) - \frac{C}{\sqrt{k}} \text{Id} \\ &= \sum_{\gamma \in \Gamma} J_\gamma^{(k)} L_\gamma (H_0^{(k)} + M^{(k)} E_\lambda^{(k)}) L_\gamma^{-1} J_\gamma^{(k)} - \frac{C}{\sqrt{k}} \text{Id} \\ &\geq \sum_{\gamma \in \Gamma} J_\gamma^{(k)} L_\gamma \lambda L_\gamma^{-1} J_\gamma^{(k)} - \frac{C}{\sqrt{k}} \text{Id} \\ &= \left(\lambda - \frac{C}{\sqrt{k}} \right) \text{Id}. \end{aligned} \quad (11)$$

Lemma 2.5. $\text{rank}_\Gamma G^{(k)} \leq N(\lambda, H_0^{(k)})$.

Proof. The operator

$$L_\alpha^{(1)} : \bigoplus_{\gamma \in \Gamma} L^2(U_{s,\gamma}, V) \longrightarrow \bigoplus_{\gamma \in \Gamma} L^2(U_{s,\gamma}, V),$$

$$L_\alpha^{(1)}((w_\gamma)_\gamma) = (w_{\alpha^{-1}\gamma})_\gamma$$

is a unitary operator for any $\alpha \in \Gamma$. Consider $i : L^2(M, V) \rightarrow \bigoplus_{\gamma \in \Gamma} L^2(U_{s,\gamma}, V)$, $i(u) = (u|_{U_{s,\gamma}})_\gamma$. Then $\|u\| \leq \|i(u)\| \leq C_1\|u\|$ and hence i is into and bounded. Moreover $L_\alpha^{(1)}i = iL_\alpha$, for $\alpha \in \Gamma$. Let

$$F : \bigoplus_{\gamma \in \Gamma} L^2(U_{s,\gamma}, V) \rightarrow L^2(M, V),$$

$$F((w_\gamma)_\gamma) = \sum_{\gamma \in \Gamma} w_\gamma.$$

F is onto, bounded and $F L_\alpha^{(1)} = L_\alpha F$, $\alpha \in \Gamma$. We define

$$\tilde{G}^{(k)} : \bigoplus_{\gamma \in \Gamma} L^2(U_{s,\gamma}, V) \rightarrow \bigoplus_{\gamma \in \Gamma} L^2(U_{s,\gamma}, V),$$

$$\tilde{G}^{(k)}((w_\gamma)_\gamma) = (J_\gamma^{(k)} L_\gamma^{-1} M^{(k)} E_\lambda^{(k)} L_\gamma J_\gamma^{(k)} w_\gamma)_\gamma.$$

Then $\tilde{G}^{(k)}$ is bounded, commutes with $L_\alpha^{(1)}$ and $G^{(k)} = F \tilde{G}^{(k)} i$. We define also the operator

$$K : \bigoplus_{\gamma \in \Gamma} L^2(U_{s,e}, V) \rightarrow \bigoplus_{\gamma \in \Gamma} L^2(U_{s,\gamma}, V),$$

$$K((w_\gamma)_\gamma) = (L_\gamma w_\gamma)_\gamma.$$

which is unitary and $K L_\alpha^{(2)} = L_\alpha^{(1)} K$, $\alpha \in \Gamma$ where

$$L_\alpha^{(2)} : \bigoplus_{\gamma \in \Gamma} L^2(U_{s,e}, V) \rightarrow \bigoplus_{\gamma \in \Gamma} L^2(U_{s,e}, V),$$

$$L_\alpha^{(2)}((w_\gamma)_\gamma) = (w_{\alpha^{-1}\gamma})_\gamma.$$

Finally, let

$$\bar{G}^{(k)} : \bigoplus_{\gamma \in \Gamma} L^2(U_{s,e}, V) \rightarrow \bigoplus_{\gamma \in \Gamma} L^2(U_{s,e}, V),$$

$$\bar{G}^{(k)}((w_\gamma)_\gamma) = (J_e^{(k)} M^{(k)} E_\lambda^{(k)} J_e^{(k)} w_\gamma)_\gamma.$$

Then

$$K \bar{G}^{(k)} = \tilde{G}^{(k)} K. \quad (12)$$

As $G^{(k)} = F \tilde{G}^{(k)} i$, we have that $\text{rank}_\Gamma G^{(k)} \leq \text{rank}_\Gamma \tilde{G}^{(k)}$. The operator K being unitary it follows from (12) that $\text{rank}_\Gamma \tilde{G}^{(k)} = \text{rank}_\Gamma \bar{G}^{(k)}$. But $R(\bar{G}^{(k)})$ is closed because $R(\bar{G}_e^{(k)})$ is closed ($\bar{G}_e^{(k)}$ is the component of $\bar{G}^{(k)}$ on $L^2(U_{s,e}, V)$ and has finite rank). If we identify $\bigoplus_{\gamma \in \Gamma} L^2(U_{s,e}, V)$ with $L^2\Gamma \otimes L^2(U_{s,e}, V)$ and consider $\bar{G}^{(k)}$ as an operator in $L^2\Gamma \otimes L^2(U_{s,e}, V)$, then $R(\bar{G}^{(k)})$ corresponds to $L^2\Gamma \otimes R(\bar{G}_e^{(k)})$ and hence

$$\text{rank}_\Gamma \bar{G}^{(k)} = \text{rank} \bar{G}_e^{(k)} \leq \text{rank} E_\lambda^{(k)} = N(\lambda, H_0^{(k)}).$$

Now the conclusion follows from the inequality

$$\text{rank}_\Gamma G^{(k)} \leq \text{rank}_\Gamma \tilde{G}^{(k)} = \text{rank}_\Gamma \bar{G}^{(k)}.$$

□

Proposition 2.6. *There is a constant $C \geq 0$ such that*

$$N_\Gamma \left(\lambda, \frac{1}{k} \Delta''_{k,q} \right) \leq N \left(\lambda + \frac{C}{\sqrt{k}}, \frac{1}{k} \Delta''_{k,q} \mid U_{k^{-\frac{1}{4}}} \right) \quad \lambda \in \mathbb{R}, k \in \mathbb{N}^* \quad (13)$$

Proof. Proposition 2.3 with $\mu = \lambda - \frac{C}{\sqrt{k}}$ and $p = N \left(\lambda, \frac{1}{k} \Delta''_{k,q} \mid U_{k^{-\frac{1}{4}}} \right)$, (11) and Lemma 2.5 entail

$$N_\Gamma \left(\lambda - \frac{C}{\sqrt{k}} - \varepsilon, \frac{1}{k} \Delta''_{k,q} \right) \leq N \left(\lambda, \frac{1}{k} \Delta''_{k,q} \mid U_{\frac{1}{\sqrt[4]{k}}} \right), \varepsilon > 0$$

Replacing λ with $\lambda + \frac{C}{\sqrt{k}} + \varepsilon$, we obtain

$$N_\Gamma \left(\lambda, \frac{1}{k} \Delta''_{k,q} \right) \leq N \left(\lambda + \frac{C}{\sqrt{k}} + \varepsilon, \frac{1}{k} \Delta''_{k,q} \mid U_{k^{-\frac{1}{4}}} \right)$$

When $\varepsilon \rightarrow 0$ it follows

$$N_\Gamma \left(\lambda, \frac{1}{k} \Delta''_{k,q} \right) \leq N \left(\lambda + \frac{C}{\sqrt{k}}, \frac{1}{k} \Delta''_{k,q} \mid U_{k^{-\frac{1}{4}}} \right)$$

□

3 Holomorphic Morse Inequalities

Let M be a Riemannian manifold of dimension n with volume element $d\sigma$. Let E and F be hermitian vector bundles on M , $\text{rank } E = 1$, $\text{rank } F = r$, with D and ∇ the canonical connections, S a continuous section in $\Lambda_{\mathbb{R}}^1 T^* M \otimes_{\mathbb{R}} \text{Hom}_{\mathbb{C}}(F, F)$ and V a continuous section in $\text{Herm}(F)$. Let ∇_k be the connection in $E^k \otimes F$. We denote the endomorphisms $\text{Id}_{E^k} \otimes S$ and $\text{Id}_{E^k} \otimes V$ by S and V . Given $\Omega \Subset M$, let

$$Q_{\Omega,k}(u) = \int_{\Omega} \left(\frac{1}{k} \mid \nabla_k u + Su \mid^2 - (Vu, u) \right) d\sigma, \quad (14)$$

$$\text{Dom}(Q_{\Omega,k}) = W_0^1(\Omega, E^k \otimes F)$$

where by W_0^1 we denote the Sobolev space. Let $V_1(x) \leq \dots \leq V_r(x)$ be the eigenvalues of $V(x)$. We shall use the following

Theorem 3.1 (Demailly [4]). *The counting function of the eigenvalues of $Q_{\Omega,k}$ satisfies for every $\lambda \in \mathbb{R}$ the following asymptotic estimates as $k \rightarrow \infty$:*

$$\sum_{j=1}^r \int_{\Omega} \nu_B(V_j + \lambda) d\sigma \leq \liminf k^{-\frac{n}{2}} N(\lambda, Q_{\Omega,k}) \leq \limsup k^{-\frac{n}{2}} N(\lambda, Q_{\Omega,k}) \leq \sum_{j=1}^r \int_{\overline{\Omega}} \bar{\nu}_B(V_j + \lambda) d\sigma \quad (15)$$

where B is the magnetic field of the connection D and

$$\nu_B(\lambda) = \frac{2^{s-n} \pi^{-\frac{n}{2}}}{\Gamma\left(\frac{n}{2} - s + 1\right)} B_1 \cdots B_s \sum_{(p_1, \dots, p_n) \in \mathbb{N}^s} \left[\lambda - \sum_{j=1}^s (2p_j + 1) B_j \right]_+^{\frac{n}{2} - s} \quad (16)$$

if $B_1(x) \geq \dots \geq B_s(x)$ are the absolute values of the non-zero eigenvalues of B , $[\lambda]_+^0 = 0$ for $\lambda \leq 0$, $[\lambda]_+^0 = 1$ for $\lambda > 0$ and $\bar{\nu}_B = \lim \nu_B(\lambda + \varepsilon), \varepsilon \searrow 0$.

We assume now that M is a hermitian complex analytic manifold of complex dimension n , E and F hermitian holomorphic vector bundles, $F_k = E^k \otimes F$ and Δ_k'' the Laplace–Beltrami operator on F_k . If $\alpha_1(x), \dots, \alpha_n(x)$ are the eigenvalues of $ic(E)(x)$ with respect to the metric on M then, as in Demailly [4], we deduce that there is a countable set $A \subset \mathbb{R}$ such that

$$\lim k^{-n} N \left(\lambda, \frac{1}{k} \Delta_{k,q}'' \mid \Omega \right) = r \sum_{|J|=q} \int_{\Omega} \nu_B (2\lambda + \alpha_{C(J)} - \alpha_J) d\sigma \quad (17)$$

for $\lambda \in \mathbb{R} \setminus A$, where $\alpha_J = \sum_{j \in J} \alpha_j$, $C(J) = \{1, \dots, n\} \setminus J$.

Let M be an analytic complex manifold of dimension n and Γ a discrete group which acts freely and properly discontinuous on M such that $X = M/\Gamma$ is compact. We choose a hermitian metric on X and we lift it on M . Let E and F be hermitian holomorphic vector bundles on X , $\text{rank } E = 1$, $\text{rank } F = r$ and $\tilde{E} = \pi^* E$, $\tilde{F} = \pi^* F$. Let $E(\cdot, \Delta_{k,q}'')$ the spectral family of the self-adjoint operator $\Delta_{k,q}''$ in $L_{0,q}^2(M, \tilde{E}^k \otimes \tilde{F})$ and $L_q^{\lambda,k} = R \left(E \left([0, \lambda], \frac{1}{k} \Delta_{k,q}'' \right) \right)$. Then

$$E([0, \lambda], \Delta_{k,q}'') \bar{\partial}_{q-1} = \bar{\partial}_{q-1} E([0, \lambda], \Delta_{k,q}'')$$

on $L_{q-1}^{\lambda,k}$ and it follows that $\bar{\partial}_{q-1} L_{q-1}^{\lambda,k} \subset L_q^{\lambda,k}$. If $\bar{\partial}_q^\lambda$ denotes the restriction

$$\bar{\partial}_q : L_q^{\lambda,k} \rightarrow L_{q+1}^{\lambda,k}$$

then

$$\{u \in L_q^{\lambda,k} \mid \bar{\partial}_q^\lambda u = 0, (\bar{\partial}_{q-1}^\lambda)^* u = 0\} = \mathcal{H}_{(2)}^q(M, \tilde{E}^k \otimes \tilde{F}) \quad (18)$$

(see Shubin [8]). By definition $N_r(\lambda, \frac{1}{k} \Delta_{k,q}'') = \dim_r L_q^{\lambda,k}$. There is a complex

$$0 \rightarrow L_0^{\lambda,k} \rightarrow L_1^{\lambda,k} \rightarrow \dots \rightarrow L_n^{\lambda,k} \rightarrow 0 \quad (19)$$

From Proposition 2.1 we get

$$\sum_{j=1}^q (-1)^{q-j} \dim_r (N(\bar{\partial}_q^\lambda) / \overline{R(\bar{\partial}_{q-1}^\lambda)}) \leq \sum_{j=1}^q (-1)^q N_r \left(\lambda, \frac{1}{k} \Delta_{k,q}'' \right)$$

for $q = 0, 1, \dots, n$ and for $q = n$ the inequality becomes equality.

From (18), $N(\bar{\partial}_q^\lambda) / \overline{R(\bar{\partial}_{q-1}^\lambda)} \simeq \mathcal{H}_{(2)}^q(M, \tilde{E}^k \otimes \tilde{F})$. From Theorem 2.4 and Proposition 2.6 it follows that there is a constant C such that

$$N(\lambda, \frac{1}{k} \Delta_{k,q}'') \mid U \leq N_r(\lambda, \frac{1}{k} \Delta_{k,q}'') \leq N \left(\lambda + \frac{C}{\sqrt{k}}, \frac{1}{k} \Delta_{k,q}'' \mid U_{\frac{1}{\sqrt{k}}} \right) \quad (20)$$

But

$$N(\lambda, \frac{1}{k} \Delta_{k,q}'') \mid U = r k^n \sum_{|J|=q} \int_U \nu_B (2\lambda + \alpha_{C(J)} - \alpha_J) d\sigma + o(k^n)$$

and

$$\begin{aligned} \limsup k^{-n} N \left(\lambda + \frac{C}{\sqrt{k}}, \frac{1}{k} \Delta_{k,q}'' \mid U_{\frac{1}{\sqrt{k}}} \right) &\leq \lim k^{-n} N(\lambda + \varepsilon, \frac{1}{k} \Delta_{k,q}'') \mid U_\varepsilon \\ &= r \sum_{|J|=q} \int_{U_\varepsilon} \nu_B (2\lambda + 2\varepsilon + \alpha_{C(J)} - \alpha_J) d\sigma \end{aligned}$$

so when $\varepsilon \rightarrow 0$ we get

$$\lim k^{-n} N \left(\lambda + \frac{C}{\sqrt{k}}, \frac{1}{k} \Delta''_{k,q} \mid U_{k^{-\frac{1}{4}}} \right) = r \sum_{|J|=q} \int_{\bar{U}} \bar{\nu}_B (2\lambda + \alpha_{C(J)} - \alpha_J) d\sigma$$

for every $\lambda \in \mathbb{R} \setminus A$. As $\partial U = \bar{U} \setminus U$ is of measure zero because U is a fundamental domain, it follows that

$$N_\Gamma(\lambda, \frac{1}{k} \Delta''_{k,q}) = r k^n \sum_{|J|=q} \int_U \nu_B (2\lambda + \alpha_{C(J)} - \alpha_J) d\sigma + o(k^n)$$

for $\lambda \in \mathbb{R} \setminus A$. Hence for $\lambda \rightarrow 0$, $\lambda \in \mathbb{R} \setminus A$ we obtain

$$\sum_{j=0}^q (-1)^{q-j} \dim_\Gamma \mathcal{H}_{(2)}^j(M, \tilde{E}^k \otimes \tilde{F}) \leq k^n \sum_{j=0}^q (-1)^{q-j} I^j + o(k^n) \quad (21)$$

where

$$I^j = r \sum_{|J|=q} \int_U \nu_B (\alpha_{C(J)} - \alpha_J) d\sigma$$

which is calculated in Demailly [4] :

$$I^j = \frac{r}{n!} \int_{M(j) \cap U} (-1)^j \left(\frac{i}{2\pi} c(\tilde{E}) \right)^n = \frac{r}{n!} \int_{X(j)} (-1)^j \left(\frac{i}{2\pi} c(E) \right)^n \quad (22)$$

with $M(j) = \{x \in M \mid ic(\tilde{E})(x) \text{ has } j \text{ negative eigenvalues and } n-j \text{ positive ones}\}$. We have used that $c(\tilde{E})$ is the lifting of $c(E)$. Theorem 1.1 now follows from (21) and (22).

References

- [1] A. Andreotti, E. Vesentini, *Carleman estimates for the Laplace-Beltrami equation on complex manifolds*, Inst. Hautes Etudes Sci., Publ. Math. No. **25** (1965), 81–130.
- [2] M. F. Atiyah, *Elliptic operators, discrete groups and von Neumann algebras*, Asterisque, **32–33** (1976), 43–72.
- [3] T. Bouche, *Inegalités de Morse pour la d'' -cohomologie sur une variété non-compacte*, Ann.Sci. Ecole Norm.Sup. **22** (1989), 501–513.
- [4] J. P. Demailly, *Champs magnétiques et inegalités de Morse pour la d'' -cohomologie*, Ann. Inst. Fourier **35**(4) (1985), 189–229.
- [5] J. Kollár, *Shafarevich maps and automorphic forms*, Princeton University Press, Princeton, NJ, 1995.
- [6] G. Marinescu, *Asymptotic Morse Inequalities for Pseudoconcave Manifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **23** (1996), no. 1, 27–55.
- [7] T. Napier, *Convexity properties of coverings of smooth projective varieties*, Math. Ann., **286** (1990), 433–479.
- [8] M. Shubin, *Semiclassical asymptotics on covering manifolds and Morse inequalities*, Geom. Funct. Anal. **6** (1996), no. 2, 370–409.

Address of the authors: Department of Mathematics,
University of Bucharest, Str. Academiei 14, 70109 Bucharest, Romania.
e-mail: radone@skylab.math.unibuc.ro, chiose@skylab.math.unibuc.ro